Catalog #: PHH1259



Summary

Name PACSIN2

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Protein Kinase C and Casein Kinase Substrate in

> Neurons Protein 2 is produced by our Mammalian expression system and the target gene encoding Met1-Gln486 is expressed with a 6His tag at the C-

terminus.

Accession # O9UNF0

Host **Human Cells**

Species Human

Predicted Molecular Mass 56.7 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

The product is shipped at ambient temperature. Upon receipt, store it **Shipping**

immediately at the temperature listed below.


Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human PACSIN2 (C-6His) Catalog #: PHH1259



Background

Alternative Names Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2; PACSIN2

Background Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2 (PACSIN2) is a

member of the PACSIN family. PACSIN2 is localized to the plasma membrane via its coiled-coil domain. PACSIN2 is widely expressed and contains one FCH domain and one SH3 domain. PACSIN2 forms homo- and hetero-aggregates with other PACSINs. PACSIN2 may play a role in vesicle formation and transport. In addition, PACSIN2 is involved in linking the actin cytoskeleton with vesicle formation by

regulating tubulin polymerization.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838