

Product Name: Recombinant Human PTH (N-8His)
Catalog #: PHH1264




Summary

Name	Parathyroid Hormone/PTH
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Parathyroid Hormone is produced by our Mammalian expression system and the target gene encoding Ser32-Gln115 is expressed with a 8His tag at the N-terminus.
Accession #	P01270
Host	Human Cells
Species	Human
Predicted Molecular Mass	10.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human PTH (N-8His)
Catalog #: PHH1264



Alternative Names

Parathyroid hormone; PTH; Parathormone; Parathyrin; PTH

Background

Parathyroid hormone (PTH) is a critical hormone in the regulation of Ca^{++} homeostasis. Parathyroid hormone is the most important endocrine regulator of calcium and phosphorus concentration in extracellular fluid. This hormone is secreted from cells of the parathyroid glands and finds its major target cells in bone and kidney. Another hormone, parathyroid hormone-related protein, binds to the same receptor as parathyroid hormone and has major effects on development. Like most other protein hormones, parathyroid hormone is synthesized as a preprohormone. After intracellular processing, the mature hormone is packaged with in the Golgi into secretory vesicles, the secreted into blood by exocytosis. In renal epithelium, PTH promotes conversion of Vitamin D to its active form, lowers Ca^{++} excretion and increases phosphate excretion. PTH also increases hematopoietic stem cell proliferation and mobilization and induces arterial vasodilation by regulating Ca^{++} influx in PTH1R-expressing arterial smooth muscle.

Note

For Research Use Only , Not for Diagnostic Use.