

**Product Name: Recombinant Human PCDH1 (C-6His)**  
**Catalog #: PHH1270**




## Summary

<b>Name</b>	PCDH1/Protocadherin-1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Protocadherin-1 is produced by our Mammalian expression system and the target gene encoding Thr58-Asn852 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	Q08174
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	87.3 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human PCDH1 (C-6His)**  
**Catalog #: PHH1270**



### Alternative Names

Protocadherin-1; Cadherin-Like Protein 1; Protocadherin-42; PC42; PCDH1

### Background

Protocadherin-1, also known as Cadherin-like protein 1, Protocadherin-42 and PCDH1, belongs to the protocadherin subfamily within the cadherin superfamily. PCDH1 contains seven cadherin-like domains, a transmembrane region and a C-terminal cytoplasmic region. PCDH1 can be detected as early as embryonic day 9.5. In early embryogenesis, expression is especially prominent in blood vessels. The tight spatial and temporal regulation of Pcdh1 expression suggests that this protocadherin plays multiple roles not only during development but also in mature tissues and organs. In addition, protocadherin-1 is involved in cell-cell interaction processes and in neural cell adhesion.

### Note

For Research Use Only , Not for Diagnostic Use.