

Summary

Name	PDGF R alpha/PDGFRA/CD140a
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Platelet-derived Growth Factor Receptor Alpha is produced by our Mammalian expression system and the target gene encoding Gln24-Glu524 is expressed with a 6His tag at the C-terminus.
Accession #	P16234
Host	Human Cells
Species	Human
Predicted Molecular Mass	57 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human PDGFRA (C-6His)
Catalog #: PHH1283



Alternative Names

Platelet-derived growth factor receptor alpha; PDGFR-alpha; Alpha platelet-derived growth factor receptor; CD140 antigen-like family member A; Platelet-derived growth factor alpha receptor; Platelet-derived growth factor receptor 2; PDGFR-2; CD140a

Background

Platelet-derived Growth Factor Receptor Alpha (PDGF $R\alpha$) is an enzyme that belongs to the class III subfamily of receptor tyrosine kinases. It is a type I transmembrane glycoprotein, and can form homo- or hetero-dimeric receptors when engaged by dimers of the PDGF family of growth factors. PDGF $R\alpha$ is strongly expressed in oligodendrocyte, lung, skin and intestinal progenitor cells and induced by inflammation or growth in culture, but is lowly expressed in most mesenchymal cells. PDGF $R\alpha$ autophosphorylates upon dimerization, activating signaling cascades in PI-3kinase Ras-MAP kinase, and PLC- γ pathways. PDGF $R\alpha$ has influence on local gradients of epithelially produced PDGF-AA or PDGF-CC during formation of the cranial, cardiac neural crest and interstitial kidney mesenchyme.

Note

For Research Use Only , Not for Diagnostic Use.