Catalog #: PHH1307



Summary

Name Periostin(POSTN)

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Osteoblast Specific Factor 2 is produced by our

Mammalian expression system and the target gene encoding Asn22-Gln779

is expressed with a 6His tag at the C-terminus.

Accession # Q15063-2

Host **Human Cells**

Species Human

Predicted Molecular Mass 85.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.0.

The product is shipped at ambient temperature. Upon receipt, store it **Shipping**

immediately at the temperature listed below.

Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.


Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Alternative Names

Osteoblast-specific factor 2; OSF-2; POSTN; Periostin

Background

Osteoblast-Specific Factor OSF-2 (POSTN, Periostin, OSF-2) is a secreted, homodimeric protein that belongs to the periostin family of the FAS1 superfamily of molecules. Periostin is a disulfide linked 90kDa bone adhesion protein secreted by osteoblasts and osteoblast-like cell lines and the protein is an attachment agent for osteoblasts. It is a TGF-beta inducible molecule that serves as both an adhesion molecule and tumor suppressor. It is synthesized by smooth muscle cells, fibroblasts and osteoblasts, as well as in the periosteum and periodontal ligament. Periostin functions as a ligand for alpha-V/beta-3 and alpha-V/beta-5 integrins to support adhesion and migration of epithelial cells.

Note

For Research Use Only, Not for Diagnostic Use.