


Product Name: Recombinant Human PCSK9 (D374Y,C-6His)
Catalog #: PHH1378

Summary

Name	PCSK9/Proprotein Convertase 9 (Val474Ile,Gly670Glu,D374Y)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Proprotein Convertase Subtilisin/Kexin Type 9 is produced by our Mammalian expression system and the target gene encoding Gln31-Gln692 (Asp374Tyr,Val474Ile, Gly504Arg, Gly670Glu) is expressed with a 6His tag at the C-terminus.
Accession #	Q8NBP7
Host	Human Cells
Species	Human
Predicted Molecular Mass	15-18&60-80 KDa
Formulation	Supplied as a 0.2 μm filtered solution of 50mM HEPES, 150mM NaCl, 20% Glycerol, pH 7.4.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image

Product Name: Recombinant Human PCSK9 (D374Y,C-6His)
Catalog #: PHH1378



Alternative Names

Proprotein Convertase Subtilisin/Kexin Type 9; Neural Apoptosis-Regulated Convertase 1; NARC-1; Proprotein Convertase 9; PC9; Subtilisin/Kexin-Like Protease PC9; PCSK9; NARC1

Background

Recombinant Human Proprotein Convertase Subtilisin/Kexin Type 9/PCSK9 (D374Y) is a gain of function mutant of human PCSK9 protein. Human PCSK9 is a secretory subtilase belonging to the proteinase K subfamily. PCSK9 is synthesized as a soluble zymogen that undergoes autocatalytic intramolecular processing in the ER, the pro domain and mature chain are secreted together through noncovalent interactions. PCSK9 binds with low-density lipoprotein receptor (LDLR) and it plays a major regulatory role in cholesterol homeostasis. Inhibition of PCSK9 function by preventing PCSK9/LDLR interaction is currently being explored as a means of lowering cholesterol levels. PCSK9 also binds to apolipoprotein receptor 2 (ApoER2), and play a role in the neural development.

Note

For Research Use Only , Not for Diagnostic Use.