


Product Name: Recombinant Human PSG9 (C-6His)
Catalog #: PHH1398



Summary

Name	PSG9/PSG11
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Pregnancy-specific Beta-1-glycoprotein 9 is produced by our Mammalian expression system and the target gene encoding Glu35-Ser426 is expressed with a 6His tag at the C-terminus.
Accession #	AAH20759.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	45.6 KDa
Formulation	Supplied as a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Product Name: Recombinant Human PSG9 (C-6His)
Catalog #: PHH1398



Alternative Names

Pregnancy-specific beta-1-glycoprotein 9; PS-beta-G-9; PSBG-9; Pregnancy-specific glycoprotein 9; PS34; Pregnancy-specific beta-1 glycoprotein B; PS-beta-B; Pregnancy-specific beta-1-glycoprotein 11; PS-beta-G-11; PSBG-11; Pregnancy-specific glycoprotein 11; Pregnancy-specific glycoprotein 7; PSG7; PSG11

Background

Pregnancy-specific beta-1-glycoprotein 9 (PSG9) is a secreted protein and contains 3 Ig-like C2-type (immunoglobulin-like) domains, 1 Ig-like V-type (immunoglobulin-like) domain. It is a member of the PSG family, a group of closely related secreted glycoproteins that are highly expressed in fetal placental syncytiotrophoblast cells. The members of the PSG protein family all have a characteristic N-terminal domain that is homologous to the immunoglobulin variable region. PSGs become detectable in serum during the first two to three weeks of pregnancy and increase as the pregnancy progresses, eventually representing the most abundant fetal protein in the maternal blood at term. PSGs function to stimulate secretion of TH2-type cytokines from monocytes, and they may also modulate the maternal immune system during pregnancy, thereby protecting the semi-allogenic fetus from rejection. PSGs are commonly expressed in trophoblast tumors. Eleven human PSG proteins (PSG1-PSG11) have been described.

Note

For Research Use Only , Not for Diagnostic Use.