**Product Name: Recombinant Human Renin (C-10His)** Catalog #: PHH1424



### **Summary**

Name Renin/Angiotensinogenase/angiotensin-forming enzyme

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/µg as determined by LAL test.

Construction Recombinant Human Renin is produced by our Mammalian expression

system and the target gene encoding Leu24-Arg406 is expressed with a 10His

tag at the C-terminus.

P00797 Accession #

Host **Human Cells** 

**Species** Human

**Predicted Molecular Mass** 44 KDa

**Formulation** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 8% Sucrose, 5% Mannitol,

0.05% Tween80, 100mM NaCl, pH 7.4.

The product is shipped at ambient temperature. Upon receipt, store it **Shipping** 

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at  $\leq$  -20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at  $\leq$  -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

> not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. /xa0Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

/xa0


### **SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: Recombinant Human Renin (C-10His)**

Catalog #: PHH1424





### **Alternative Names**

Renin; Angiotensinogenase; REN

## **Background**

Renin is a member of the aspartyl proteinase family produced largely in part by the juxtaglomerular cells in the kidney. Renin is produced as prorenin with 43 pro residues at the N-terminal of mature Renin. The inactive prorenin becomes activated proteolytically by trypsin, cathepsin B, or other proteinases. Renin also has a very high selectivity for substrates due to a long peptide recognition on either side of the peptide bond undergoing cleavage. An octapeptide substrate was the minimum length to be cleaved by Renin. Renin plays a crucial role in the regulation of blood pressure and salt balance through the cleavage of angiotensinogen, which is the only known physiological substrate of Renin. Renin releases the decapeptide angiotensin I, which in turn is further converted to vasoactive hormone angiotensin II by angiotensin converting enzyme (ACE).

### Note

For Research Use Only, Not for Diagnostic Use.