


**Product Name: Recombinant Human RSPO3 (C-Fc-6His)**  
**Catalog #: PHH1449**



## Summary

<b>Name</b>	R-spondin-3/RSPO3/Rspondin-1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human R-spondin-3 is produced by our Mammalian expression system and the target gene encoding Gln22-Val201 is expressed with a human IgG1 Fc, 6His tag at the C-terminus.
<b>Accession #</b>	Q9BXY4
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	47.9 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human RSPO3 (C-Fc-6His)**  
**Catalog #: PHH1449**



---

**Alternative Names**

R-spondin-3; RSPO3; Protein with TSP type-1 repeat; Roof plate-specific spondin-3; Thrombospondin type-1 domain-containing protein 2; PWTSR; THSD2; CRISTIN1

**Background**

R-spondin-3 (RSPO3), also known as Protein with TSP type-1 repeat, Roof plate-specific spondin-3, Thrombospondin type-1 domain-containing protein 2, PWTSR, THSD2 and CRISTIN1, is a member of the thrombospondin type 1 repeat supergene family. RSPO3 is a secreted protein and widely expressed in many tissues. RSPO3 contains two Furin-like repeats which have been found in a variety of eukaryotic proteins involved in the mechanism of signal transduction by receptor tyrosine kinases, and one TSP type-1 domain, RSPO3 functions as an activator of the beta-catenin signaling cascade, leading to TCF-dependent gene activation. Otherwise, RSPO3 may negatively regulate the TGF-beta pathway.

**Note**

For Research Use Only , Not for Diagnostic Use.