

Product Name: Recombinant Human sCD4 (C-Fc)
Catalog #: PHH1466




Summary

Name	sCD4/T-cell surface glycoprotein CD4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human T-cell Surface Glycoprotein CD4 is produced by our Mammalian expression system and the target gene encoding Lys26-Trp390 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P01730
Host	Human Cells
Species	Human
Predicted Molecular Mass	67.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 1mM EDTA, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human sCD4 (C-Fc)
Catalog #: PHH1466



Alternative Names

T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; CD4; Scd4

Background

CD4 is an approximately 55 kDa type I transmembrane glycoprotein that is expressed predominantly on thymocytes and a subset of mature T lymphocytes. It is a standard phenotype marker for the identification of T cell populations. Mature human CD4 consists of a 371 amino acid extracellular region containing four immunoglobulin-like domains, a 22 amino acid transmembrane segment, and a 40 amino acid cytoplasmic domain. CD4 is expressed along with CD8 on double positive T cells during their development in the thymus. CD4 binds directly to MHC class II molecules on antigen presenting cells (10). This interaction contributes to the formation of the immunological synapse which is focused around the TCR-MHC class II-antigenic peptide interaction. CD4 also functions as a chemotactic receptor for IL-16 and, in human, as a co-receptor for the gp120 surface glycoprotein of HIV-1.

Note

For Research Use Only , Not for Diagnostic Use.