Product Name: Recombinant Human STX7 (N-6His)

Catalog #: PHH1597



Summary

Name Syntaxin-7/STX7

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Syntaxin-7 is produced by our Mammalian expression

system and the target gene encoding Ser2-Leu238 is expressed with a 6His

tag at the N-terminus.

Accession # O15400

Host Human Cells

Species Human

Predicted Molecular Mass 28 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human STX7 (N-6His)

Catalog #: PHH1597





Alternative Names

Syntaxin-7; STX7; syntaxin 7

Background

Syntaxin 7 (STX7) is a member of a family of membrane-trafficking proteins named as soluble N-ethylmaleimide-sensitive factor attachment protein receptors (SNAREs), and mediates the endocytic trafficking from early endosomes to late endosomes and lysosomes. Syntaxin-7 has been shown to be present in both late endosomes and lysosomes, and to be required for both homotypic late endosome fusion and heterotypic fusion with lysosomes. STX7 has been shown to interact with STX8, VPS18, Vesicle-associated membrane protein 8 and VPS11.

Note

For Research Use Only, Not for Diagnostic Use.