


Product Name: Recombinant Human TIGIT (C-Fc)
Catalog #: PHH1609



Summary

Name	TIGIT/VSIG9/VSTM3/T-cell immunoreceptor with Ig and ITIM domains
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human T-cell Immunoreceptor With Ig And ITIM Domains is produced by our Mammalian expression system and the target gene encoding Met22-Pro141 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q495A1
Host	Human Cells
Species	Human
Predicted Molecular Mass	39.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human TIGIT (C-Fc)
Catalog #: PHH1609



Background

Alternative Names

T-cell immunoreceptor with Ig and ITIM domains; V-set and immunoglobulin domain-containing protein 9; V-set and transmembrane domain-containing protein 3; Tigit; VSIG9; VSTM3

Background

T cell immunoreceptor with Ig and ITIM domains (TIGIT) is a member of the CD28 family within the Ig superfamily of proteins. TIGIT is expressed on NK cells and subsets of activated, memory and regulatory T cells, and particularly on follicular helper T cells within secondary lymphoid organs. It binds to CD155 and Nectin-2 that appear on dendritic cells (DC) and endothelium. Ligation of TIGIT on T cells down-regulates TCR-mediated activation and subsequent proliferation, while NK cell TIGIT ligation blocks NK cell cytotoxicity. Through CD155 and Nectin-2, which also interact with DNAM-1/CD226 and CD96/Tactile, TIGIT is part of an interacting network of Ig superfamily members that may augment or oppose each other. In particular, TIGIT binding to CD155 can antagonize the effects of DNAM1.

Note

For Research Use Only , Not for Diagnostic Use.