# **Product Name: Recombinant Human TFF2 (C-6His)**

Catalog #: PHH1618



### **Summary**

Name TFF2/Trefoil factor 2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human Trefoil Factor 2 is produced by our Mammalian

expression system and the target gene encoding Glu24-Tyr129 is expressed

with a 6His tag at the C-terminus.

Accession # Q03403

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 13 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image

# **Product Name: Recombinant Human TFF2 (C-6His)**

Catalog #: PHH1618





### **Alternative Names**

Trefoil Factor 2; Spasmolysin; Spasmolytic Polypeptide; SP; TFF2; SML1

## **Background**

Trefoil Factor 2 (TFF2) is a member of the trefoil family and contains two P-type (trefoil) domains. Members of this family are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfides. TFF2 is a secreted protein and specifically expressed in the stomach. TFF2 inhibits gastrointestinal motility and gastric acid secretion. TFF2 could function as a structural component of gastric mucus, possibly by stabilizing glycoproteins in the mucus gel through interactions with carbohydrate side chains.

#### Note

For Research Use Only, Not for Diagnostic Use.