

**Product Name: Recombinant Human TIMP-1 (C-6His)**  
**Catalog #: PHH1654**




## Summary

<b>Name</b>	TIMP-1/Metalloproteinase inhibitor 1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Tissue Inhibitor Of Metalloproteinases 1 is produced by our Mammalian expression system and the target gene encoding Cys24-Ala207 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P01033
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	21.75 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human TIMP-1 (C-6His)**  
**Catalog #: PHH1654**



### Alternative Names

Metalloproteinase Inhibitor 1; Erythroid-Potentiating Activity; EPA; Fibroblast collagenase Inhibitor; Collagenase Inhibitor; Tissue Inhibitor of Metalloproteinases 1; TIMP-1; TIMP1; CLGI; TIMP

### Background

Tissue Inhibitor of Metalloproteinases 1 (TIMP-1) is a member of TIMP family. The homologous proteins of TIMPs regulate the activity of matrix metalloproteinases (MMPs), including inhibition of active MMPs, proMMP activation, cell growth promotion, matrix binding, inhibition of angiogenesis and the induction of apoptosis. Timp-1 complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. It also mediates erythropoiesis in vitro; but, unlike IL-3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors. It is known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-11, MMP-12, MMP-13, and MMP-16, without MMP-14.

### Note

For Research Use Only , Not for Diagnostic Use.