

Product Name: Recombinant Human TIMP-2 (C-6His)
Catalog #: PHH1655




Summary

Name	TIMP-2/Metalloproteinase inhibitor 2/Tissue Inhibitor of Metalloproteinases 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Tissue Inhibitor of Metalloproteinases 2 is produced by our Mammalian expression system and the target gene encoding Cys27-Pro220 is expressed with a 6His tag at the C-terminus.
Accession #	P16035
Host	Human Cells
Species	Human
Predicted Molecular Mass	22.79 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human TIMP-2 (C-6His)
Catalog #: PHH1655



Alternative Names

Metalloproteinase Inhibitor 2; CSC-21K; Tissue Inhibitor of Metalloproteinases 2; TIMP-2; TIMP2

Background

Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). There are four members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4. Tissue Inhibitor of Metalloproteinases 2 (TIMP-2) is a non N-glycosylated protein with a molecular mass of 22 kDa. It produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-2 also has erythroid-potentiating and cell growth promoting activities.

Note

For Research Use Only , Not for Diagnostic Use.