

**Product Name: Recombinant Human TNF RII (174AA,C-6His)**  
**Catalog #: PHH1687**




---

## Summary

<b>Name</b>	TNF RII/TNFRSF1B/CD120b/TNFR2/TNF Receptor II/Tumor Necrosis Factor Receptor II
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Tumor Necrosis Factor Receptor II is produced by our Mammalian expression system and the target gene encoding Lys288-Ser461 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P20333
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	19 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human TNF RII (174AA,C-6His)**  
**Catalog #: PHH1687**



### **Alternative Names**

Tumor necrosis factor receptor superfamily member 1B; Tumor necrosis factor receptor 2; TNF-R2; Tumor necrosis factor receptor type II; TNF-RII; TNFR-II; p75; p80 TNF-alpha receptor

### **Background**

Tumor necrosis factor receptor superfamily member 1B(TNFRSF1B) is expressed by the gene TNFRSF1B. The soluble form is produced from the membrane form by proteolytic processing. It can bind to TRAF2, and interacts with BMX. It can act as the receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha.

### **Note**

For Research Use Only , Not for Diagnostic Use.