Product Name: Recombinant Human TREM-1 (C-6His) Catalog #: PHH1716



Summary

Name TREM-1/CD354/Triggering receptor expressed on myeloid cells 1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Triggering Receptor Expressed on Myeloid Cells 1 is

produced by our Mammalian expression system and the target gene

encoding Ala21-Arg200 is expressed with a 6His tag at the C-terminus.

Accession # Q9NP99

Host Human Cells

Species Human

Predicted Molecular Mass 21.3 KDa

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human TREM-1 (C-6His) Catalog #: PHH1716



Alternative Names

Triggering Receptor Expressed on Myeloid Cells 1; TREM-1; Triggering Receptor Expressed on Monocytes 1; CD354; TREM1

Background Triggering Receptor Expressed on Myelo

Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) is a transmembrane protein with a single Ig-like domain. TREM-1 associates with the adapter protein, DAP12, to deliver an activating signal. TREM-1 is expressed on blood neutrophils and monocytes, and the expression is up-regulated by bacterial LPS. TREM-1 is expressed at high levels on neutrophils of patients with microbial sepsis and in mice with a TREM-1/Fc fusion protein protected mice against LPS-induced shock.

Human TREM-1 shares 42% sequence homology with mouse TREM-1.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838