Product Name: Recombinant Human Tie-1 (C-6His) Catalog #: PHH1754

Ci EnkiLife

Summary

Name Tyrosine-protein kinase receptor Tie-1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Tyrosine Kinase With Immunoglobulin And Epidermal

Growth Factor Homology Domains 1 is produced by our Mammalian expression system and the target gene encoding Ala22-Gln760 is expressed

with a 6His tag at the C-terminus.

Accession # P35590

Host Human Cells

Species Human

Predicted Molecular Mass 81 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 500mM NaCl, 10%

Sucrose, pH8.2.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human Tie-1 (C-6His) Catalog #: PHH1754

C EnkiLife

Background

Alternative Names Tyrosine-Protein Kinase Receptor Tie-1; TIE1; TIE

Background TIE-1 (Tyrosine Kinase with Ig and EGF Homology domains 1) and TIE-2/Tek

comprise a receptor tyrosine kinase (RTK) subfamily. These receptors are expressed on endothelial and hematopoietic progenitor cells and play critical roles in angiogenesis, vasculogenesis and hematopoiesis. Human TIE-1 cDNA encodes a 1124 amino acid (aa) residue precursor protein with an 18aa signal peptide, a 727 aa extracellular domain and a 354 aa cytoplasmic domain. so far, two ligands have been described for TIE-2 [angiopoietin-1 (Ang1) and angiopoietin-2 (Ang2, but no

ligand was found for TIE-1.

Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838