


Product Name: Recombinant Human VSIG4 (C-Fc)
Catalog #: PHH1827



Summary

Name	VSIG4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human V-Set And Immunoglobulin Domain-Containing Protein 4 is produced by our Mammalian expression system and the target gene encoding Arg20-Val284 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q9Y279
Host	Human Cells
Species	Human
Predicted Molecular Mass	56.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human VSIG4 (C-Fc)
Catalog #: PHH1827



Alternative Names

V-set and immunoglobulin domain-containing protein 4; VSIG4; Protein Z39Ig; Z39IG; CRlg

Background

V-set and immunoglobulin domain-containing protein 4 (VSIG4) is a transmembrane protein contains a signal peptide, a V-type Ig-like domain, a C2-type Ig-like domain, several potential O-glycosylation sites, and an intracellular domain with 2 potential phosphorylation sites and is structurally related to the B7 family of immune regulatory proteins. This protein is also a receptor for the complement component 3 fragments C3b and iC3b. The main function is strong negative regulator of T-cell proliferation and IL2 production and it is also potent inhibitor of the alternative complement pathway convertases. It abundantly expressed in several fetal tissues such as adult tissues, highest expression in lung and placenta and it also expressed in resting macrophages.

Note

For Research Use Only , Not for Diagnostic Use.