

Product Name: Recombinant Human AZGP1 (C-6His)
Catalog #: PHH1846




Summary

Name	Zinc-alpha-2-glycoprotein/ZAG/AZGP1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Zinc-alpha-2-Glycoprotein is produced by our Mammalian expression system and the target gene encoding Gln21-Ser298 is expressed with a 6His tag at the C-terminus.
Accession #	P25311
Host	Human Cells
Species	Human
Predicted Molecular Mass	33.18 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 7.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human AZGP1 (C-6His)
Catalog #: PHH1846



Alternative Names

Zinc-Alpha-2-Glycoprotein; Zn-Alpha-2-GP; Zn-Alpha-2-Glycoprotein; AZGP1; ZAG; ZNGP1

Background

Zinc- α -2-Glycoprotein (AZGP1) can be found in blood plasma, seminal plasma, urine, sweat, saliva, liver, and epithelial cells of various human glands. AZGP1 has been proposed in the regulation of body weight, and the melanin production by normal and malignant melanocytes. AZGP1 stimulates lipid degradation in adipocytes and causes the extensive fat losses associated with some advanced cancers. AZGP1 has been reported to stimulate lipid breakdown and may have an important role in lipid homeostasis. Mature human AZGP1 consists of one MHC class I antigen region and a C2-type Ig-like domain. AZGP1 has two alternate splice forms, one shows a 66 amino acids substitution for the C-terminal 30 amino acids, the other one shows a nine Lys substitution for amino acid 151-298.

Note

For Research Use Only , Not for Diagnostic Use.