


Product Name: Recombinant Human CD157 (C-6His)
Catalog #: PHH1852



Summary

Name	CD157/BST1/ADP-ribosyl cyclase 2/Cyclic ADP-ribose hydrolase 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human ADP-ribosyl Cyclase/Cyclic ADP-ribose Hydrolase 2 is produced by our Mammalian expression system and the target gene encoding Gly29-Lys292 is expressed with a 6His tag at the C-terminus.
Accession #	Q10588
Host	Human Cells
Species	Human
Predicted Molecular Mass	30.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CD157 (C-6His)
Catalog #: PHH1852



Alternative Names

ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 2; ADP-ribosyl cyclase 2; Bone marrow stromal antigen 1; BST-1; Cyclic ADP-ribose hydrolase 2; cADPr hydrolase 2; CD157

Background

The cluster of differentiation (CD) system is a glycosyl phosphatidylinositol anchored membrane protein that belongs to the CD38 family. It is generally used in immunophenotyping. CD157 was discovered in a bone marrow stromal cell line where it facilitates pre-B-cell growth. CD157 is a bifunctional ectoenzyme that exhibits both ADP-ribosyl cyclase and cyclic ADP ribose hydrolase activities followed with CD38. It plays a role in rheumatoid arthritis (RA) due to its enhanced expression in RA-derived bone marrow stromal cell lines. Studies have shown that this protein has a role in predicted to function as a cell surface receptor and an immunoregulatory molecule.

Note

For Research Use Only , Not for Diagnostic Use.