


Product Name: Recombinant Human HGF (C-6His)
Catalog #: PHH1862



Summary

| | |
|---------------------------------|--|
| Name | HGF/Hepatopoietin-A/Hepatocyte Growth Factor |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <0.01 EU/μg as determined by LAL test. |
| Construction | Recombinant Human Hepatocyte Growth Factor is produced by our Mammalian expression system and the target gene encoding Gln32-Ser728 is expressed with a 6His tag at the C-terminus. |
| Accession # | P14210 |
| Host | Human Cells |
| Species | Human |
| Predicted Molecular Mass | 2653.7 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of 20mM Tris , 500mM NaCl, 3%Trehalose, 0.02% Tween 80, pH 8.0. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background

Product Name: Recombinant Human HGF (C-6His)
Catalog #: PHH1862



Alternative Names

Hepatocyte growth factor; HPTA; HGF; SF; Scatter factor; Hepatopoietin-A

Background

HGF, is a pleiotropic protein in the Plasminogen subfamily of S1 peptidases and contains 4 kringle domains, 1 PAN domain and 1 peptidase S1 domain. HGF is secreted as an inactive 728 amino acid (aa) single chain propeptide. It is cleaved after the fourth Kringle domain by a serine protease to form bioactive disulfide-linked HGF with a 60 kDa alpha and 30 kDa beta chain. HGF binds heparan-sulfate proteoglycans and the widely expressed receptor tyrosine kinase, HGF R/c-MET. HGF regulates epithelial morphogenesis by inducing cell scattering and branching tubulogenesis. It can also alter epithelium morphology by the induction of nectin-1 alpha ectodomain shedding, an adhesion protein component of adherens junctions. HGF regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto oncogenic c-Met receptor. HGF is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorigenesis, and tissue regeneration.

Note

For Research Use Only , Not for Diagnostic Use.