

**Product Name: Recombinant Human CD157 (C-Fc)**  
**Catalog #: PHH1879**




---

## Summary

<b>Name</b>	CD157/BST1/ADP-ribosyl cyclase 2/Cyclic ADP-ribose hydrolase 2
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human ADP-ribosyl cyclase/cyclic ADP-ribose Hydrolase 2 is produced by our Mammalian expression system and the target gene encoding Gly29/xadLys292 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	Q10588
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	57.1 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human CD157 (C-Fc)**  
**Catalog #: PHH1879**



### Alternative Names

ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 2; ADP-ribosyl cyclase 2; Bone marrow stromal antigen 1; BST-1; Cyclic ADP-ribose hydrolase 2; cADPr hydrolase 2; CD157

### Background

The cluster of differentiation (CD) system is a glycosyl phosphatidylinositol anchored membrane protein that belongs to the CD38 family. It is generally used in immunophenotyping. CD157 was discovered in a bone marrow stromal cell line where it facilitates pre-B-cell growth. CD157 is a bifunctional ectoenzyme that exhibits both ADP-ribosyl cyclase and cyclic ADP ribose hydrolase activities followed with CD38. It plays a role in rheumatoid arthritis (RA) due to its enhanced expression in RA-derived bone marrow stromal cell lines. Studies have shown that this protein has a role in predicted to function as a cell surface receptor and an immunoregulatory molecule.

### Note

For Research Use Only , Not for Diagnostic Use.