


## Summary

<b>Name</b>	Flt-3/Flk-2/CD135/Tyrosine-protein kinase receptor FLT3
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Receptor-Type Tyrosine-Protein Kinase FLT3/FLT3 is produced by our Mammalian expression system and the target gene encoding Asn27-Asn541 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	AAI26351.1
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	85.3 KDa
<b>Formulation</b>	Supplied as a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	

## SDS-PAGE image

**Product Name: Recombinant Human FLT3 (C-Fc)**  
**Catalog #: PHH1921**



### **Alternative Names**

Receptor-Type Tyrosine-Protein Kinase FLT3; FL Cytokine Receptor; Fetal Liver Kinase-2; FLK-2; Fms-Like Tyrosine Kinase 3; FLT-3; Stem Cell Tyrosine Kinase 1; STK-1; CD135; FLT3; FLK2; STK1

### **Background**

The Flt-3 (fms-like tyrosine kinase) receptor, also named Flk-2 and Stk-1 is a member of the class III subfamily of receptor tyrosine kinases that also includes KIT, the receptor for SCF and FMS, the receptor for M-CSF. The extracellular region of these receptors contains five immunoglobulin-like domains and the intracellular region contains a split kinase domain. Human Flt-3 cDNA encodes a 993 amino acid (aa) residue type I membrane protein with a 26 aa residue signal peptide, a 515 aa extracellular domain with 10 potential N-linked glycosylation sites, a 21 aa residue transmembrane domain and a 431 aa residue cytoplasmic domain. Flt-3 expression has been detected in various tissues, including placenta, gonads, and tissues of nervous and hematopoietic origin. Among hematopoietic cells, the expression of Flt-3 was found to be restricted to the highly enriched stem/progenitor cell populations. The ligand for Flt-3 (FL) has been identified to be a transmembrane protein with structural homology to M-CSF and SCF. Recombinant soluble Flt-3/Fc chimeric protein has been shown to bind FL with high affinity and is a potent FL antagonist.

### **Note**

For Research Use Only , Not for Diagnostic Use.