Product Name: Recombinant Human 4-1BB (C-mFc)

Catalog #: PHH2030



Summary

Name 4-1BB/CD137/TNFRSF9/ILA

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human 4-1BB Ligand Receptor is produced by our Mammalian

expression system and the target gene encoding Leu24-Gln186 is expressed

with a mouse IgG1 Fc tag at the C-terminus.

Accession # Q07011

Host Human Cells

Species Human

Predicted Molecular Mass 43.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human 4-1BB (C-mFc)

Catalog #: PHH2030



Alternative Names CD137; ILA; TNFRSF9; 4-1BB ligand receptor; CDw137; T-cell antigen 4-1BB

homolog; T-cell antigen ILA

Background Tumor necrosis factor receptor superfamily member 9(TNFRSF9) is an inducible T

cell surface protein belonging to the TNF receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. It is absent from naive T cells, but upregulated and continually expressed following T cell activation. It is a

receptor for TNFSF9/4-1BBL, and possibly active during T cell activation.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838