## **Product Name: Recombinant Human 4-1BB (C-mFc)**

Catalog #: PHH2030



## **Summary**

Name 4-1BB/CD137/TNFRSF9/ILA

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human 4-1BB Ligand Receptor is produced by our Mammalian

expression system and the target gene encoding Leu24-Gln186 is expressed

with a mouse IgG1 Fc tag at the C-terminus.

Accession # Q07011

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 43.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image

# **Product Name: Recombinant Human 4-1BB (C-mFc)**

Catalog #: PHH2030





#### **Alternative Names**

CD137; ILA; TNFRSF9; 4-1BB ligand receptor; CDw137; T-cell antigen 4-1BB homolog; T-cell antigen ILA

## **Background**

Tumor necrosis factor receptor superfamily member 9(TNFRSF9) is an inducible T cell surface protein belonging to the TNF receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. It is absent from naive T cells, but upregulated and continually expressed following T cell activation. It is a receptor for TNFSF9/4-1BBL, and possibly active during T cell activation.

#### Note

For Research Use Only, Not for Diagnostic Use.