


Product Name: Recombinant Human GITR (C-mFc)
Catalog #: PHH2039



Summary

Name	GITR/TNFRSF18/CD357/Tumor necrosis factor receptor superfamily member 18/Glucocorticoid-induced TNFR-related protein/AITR
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant human Glucocorticoid Induced TNF Receptor Family Related Gene is produced by our Mammalian expression system and the target gene encoding Gln26-Glu161(Thr45Ala) is expressed with a mouse IgG1 Fc tag at the C-terminus.
Accession #	Q9Y5U5
Host	Human Cells
Species	Human
Predicted Molecular Mass	41.1 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Alternative Names

Product Name: Recombinant Human GITR (C-mFc)
Catalog #: PHH2039



Tumor necrosis factor receptor superfamily member 18; TNFRSF18; Glucocorticoid-induced TNFR-related protein; CD357; TNFRSF18; AITR; GITR

Background

Tumor necrosis factor receptor superfamily member 18(Gitr) contains 3 TNFR-Cys repeats and it have four isforms. IsformA, isformB and isformC is single-pass type I membrane protein and isformD is a secreted protein. The protein is the receptor for TNFSF18. It seems to be involved in interactions between activated T-lymphocytes and endothelial cells and in the regulation of T-cell receptor-mediated cell death. It mediated NF-kappa-B activation via the TRAF2/NIK pathway. It binds to TRAF1, TRAF2, and TRAF3, but not TRAF5 and TRAF6 and binds through its C-terminus to SIVA1/SIVA. It preferentially expressed in activated T lymphocytes and up-regulated in peripheral mononuclear cells after antigen stimulation/lymphocyte activation.

Note

For Research Use Only , Not for Diagnostic Use.