


Product Name: Recombinant Human OX40L (C-mFc)
Catalog #: PHH2044



Summary

Name	OX40 Ligand/OX40L/TNFSF4/CD252/Tumor necrosis factor ligand superfamily member 4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human OX40 Ligand is produced by our Mammalian expression system and the target gene encoding Gln51-Leu183 is expressed with a mouse IgG1 Fc tag at the C-terminus.
Accession #	P23510
Host	Human Cells
Species	Human
Predicted Molecular Mass	42.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names	Tumor necrosis factor ligand superfamily member 4; Glycoprotein Gp34; OX40 ligand; OX40L; TAX transcriptionally-activated glycoprotein 1; TNFSF4; CD252; TXGP1
Background	Tumor necrosis factor ligand superfamily member 4(TNFSF4/OX40L) is a single-pass type II membrane protein.OX40L is expressed on the surface of activated B

Product Name: Recombinant Human OX40L (C-mFc)
Catalog #: PHH2044



cells, T cells, dendritic cells and endothelial cells. OX40L binds to OX40 (CD134), a member of the TNF receptor superfamily that is expressed predominantly on activated CD4+ T cells. OX40-OX40L co-stimulates signal to promote the survival and proliferation of activated CD4+ T cells and prolong the immune response. It involved in T-cell proliferation and cytokine production. Additional, it has been found association with systemic lupus erythematosus, no association with occurrence of atherosclerosis.

Note

For Research Use Only , Not for Diagnostic Use.