Product Name: Recombinant Human Siglec-9 (C-6His) Catalog #: PHH2066



Summary

Name Siglec-9/sialic acid-binding Ig-like lectin 9

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Sialic Acid-binding Ig-like Lectin 9 is produced by our

Mammalian expression system and the target gene encoding Gln18-Gly348 is

expressed with a 6His tag at the C-terminus.

Accession # AAH35365.2

Host **Human Cells**

Species Human

Predicted Molecular Mass 36.9 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, 2mM EDTA, pH 7.4. **Formulation**

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Siglec-9 (C-6His) Catalog #: PHH2066



Alternative Names Sialic acid-binding Ig-like lectin 9; Siglec-9; CDw329; Protein FOAP-9; SIGLEC9

Background Sialic acid-binding Ig-like lectin 9(Siglec 9) is expressed by peripheral blood

leukocytes (neutrophils and monocytes but not eosinophils), and found in liver, fetal liver, bone marrow, placenta, spleen and in lower levels in skeletal muscle, fetal brain and so on. It is a putative adhesion molecule that mediates sialic-acid dependent binding to cells. It also binds to alpha-2,3- or alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic

acids on the same cell surface.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838