Product Name: Recombinant Human CA125 (C-6His)

Catalog #: PHH2080



Summary

Name CA125/MUC16

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Mucin-16 is produced by our Mammalian expression

system and the target gene encoding Gly12660-Met12923 is expressed with a

6His tag at the C-terminus.

Accession # Q8WXI7

Host Human Cells

Species Human

Predicted Molecular Mass 29.2 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine,

pH 7.5.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CA125 (C-6His)

Catalog #: PHH2080



Alternative Names

CA125 ovarian cancer antigen; CA125; CA-125; CA125MUC-16; FLJ14303; MUC16; mucin 16, cell surface associated; mucin-16

Background

MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. MUC16 is over-expressed by tumor cells including ovarian and mesothelial cancers. The transmembrane form can adhere to mesothelin in the peritoneum, facilitating metastasis of ovarian cancer to the peritoneal cavity. MUC16 also binds galectin-1 on immune cells and enhances its expression on tumor cells. MUC16-expressing tumors adhere to NK cells, down-regulate CD16 and suppress NK response, which may promote immune evasion. MUC16 is also cyclically expressed in the endometrium and may contribute to immune privilege during pregnancy. In the eye, MUC16 and other mucins protect the cornea and keep it hydrated. It is altered on the conjunctival epithelium of patients with non-Sjogren dry eye syndrome.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838