


Summary

Name	CD5L/CD5 Antigen-Like/Sp alpha/AIM
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human CD5 Antigen-Like is produced by our Mammalian expression system and the target gene encoding Ser20-Gly347 is expressed with a 6His tag at the C-terminus.
Accession #	O43866
Host	Human Cells
Species	Human
Predicted Molecular Mass	37.09 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Alternative Names

CD5 Antigen-Like; CT-2; IgM-Associated Peptide; SP-Alpha; CD5L; API6

Background CD5 Antigen-Like (CD5L) is a soluble protein that belongs to group B of the scavenger receptor cysteine-rich (SRCR) superfamily and contains three SRCR domains. CD5L is a secreted glycoprotein and expressed by macrophages presentin lymphoid tissues. It binds to myelomonocytic and lymphoid cells and may play an important role in the regulation of the innate and adaptive immune systems. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gramnegative bacteria and the SRCR domain one of CD5L retains both the LPS and LTA binding activities. Furthermore, CD5L seems to play a role as an inhibitor of apoptosis.

Note

For Research Use Only, Not for Diagnostic Use.