


**Product Name: Recombinant Human CD82 (N-Fc)**  
**Catalog #: PHH2153**



## Summary

<b>Name</b>	CD82
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human CD82 Antigen is produced by our Mammalian expression system and the target gene encoding Gly103-Gln225 is expressed with a human IgG1 Fc tag at the N-terminus.
<b>Accession #</b>	P27701
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	40.3 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human CD82 (N-Fc)**  
**Catalog #: PHH2153**

---



**Alternative Names**

CD82 antigen; CD82 molecule; CD82; IA4; IA4C33 antigen; Inducible membrane protein R2; KAI1; KAI1GR15; ST6; ST6tspan-27; Suppressor of tumorigenicity 6 protein; Tetraspanin-27; TSPAN27; Tspan-27; TSPAN274F9

**Background**

CD82 antigen, also known as Kai-1, is a widely expressed palmitoylated molecule of the tetraspanin superfamily. KAI1/CD82 is localized on cell membrane and form interactions with other tetraspanins, integrins and chemokines which are respectively responsible for cell migration, adhesion and signaling. CD82/Kai-1 is a component of the promiscuous TIMP-1 interacting protein complex on the cell surface of human adenocarcinoma cells and gives insight into tumorigenic metastatic potential. CD82/Kai-1 suppresses EMT in prostate cancer cells adhered to fibronectin leading to reduced cell migration and invasiveness. CD82/Kai-1 function is important for muscle stem cell function in muscular disorders. Overexpression of CD82/Kai-1 suppresses growth, migration and invasion of oral cancer cells and may be considered as a potential therapeutic target in oral cancer.

**Note**

For Research Use Only , Not for Diagnostic Use.