

Product Name: Recombinant Human LILRB1 (C-Fc)
Catalog #: PHH2246




Summary

Name	LILRB1/ILT2/CD85j/Lir-1/LIR1/MIR7/Leukocyte Ig-Like Receptor B1/Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 1 is produced by our Mammalian expression system and the target gene encoding Gly24-His458 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	D9IDM8
Host	Human Cells
Species	Human
Predicted Molecular Mass	74.1 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human LILRB1 (C-Fc)
Catalog #: PHH2246



Alternative Names

Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 1; LIR-1; Leukocyte Immunoglobulin-Like Receptor 1; CD85 Antigen-Like Family Member J; Immunoglobulin-Like Transcript 2; ILT-2; Monocyte/Macrophage Immunoglobulin-Like Receptor 7; MIR-7; CD85j; LILRB1; ILT2; LIR1; MIR7

Background

The immunoglobulin-like transcript (ILT) family (also named leukocyte Ig-like receptors (LIR) and monocyte/macrophage Ig-like receptors (MIR)) can be activating and inhibitory immunoreceptors. ILTs are expressed on many leukocyte subsets and regulators of immune responses. ILTs share significant homology with killer cell Ig-like receptors (KIR). Except ILT-6, all ILT family members are type I transmembrane proteins having two or four extracellular Ig-like domains. ILT2 is expressed on most monocytes, dendritic cells, and mature B cells. ILT2 is also expressed on small percentages of T-cells and NK cells. ILT2 can prevent cellular activation.

Note

For Research Use Only, Not for Diagnostic Use.