

Product Name: Recombinant Human TRAIL R1 (C-6His)
Catalog #: PHH2280




Summary

Name	TRAIL R1/DR4/TNFRSF10A
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human TRAIL R1 is produced by our Mammalian expression system and the target gene encoding Ala24-Asn239 is expressed with a 6His tag at the C-terminus.
Accession #	O00220
Host	Human Cells
Species	Human
Predicted Molecular Mass	23.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human TRAIL R1 (C-6His)
Catalog #: PHH2280



Alternative Names

APO2; CD261 antigen; CD261; cytotoxic TRAIL receptor; DR4; TNF-related apoptosis-inducing ligand receptor 1; TNFRSF10A; TRAIL-R1; TRAILR1

Background

Tumor necrosis factor receptor superfamily member 10A (TNFRSF10A) is also known as TNF-related apoptosis-inducing ligand receptor 1 (TRAIL-R1), Death receptor 4 (DR4), CD261 and APO2, which belongs to TNF superfamily. TNFRSF10A / DR4 is widely expressed and high levels are found in spleen, peripheral blood leukocytes, small intestine and thymus, but also in K-562 erythroleukemia cells, MCF-7 breast carcinoma cells and activated T-cells. APO2 / TNFRSF10A is receptor for the cytotoxic ligand TNFSF10 / TRAIL. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF1/TRAIL), and thus transduces cell death signal and induces cell apoptosis. TRAIL R1 can promote the activation of NF-kappa-B. TRAIL R1/CD261/TNFRSF1A induces apoptosis of many transformed cell lines but not of normal tissues, even though its death domain-containing receptor, DR4, is expressed on both cell types.

Note

For Research Use Only , Not for Diagnostic Use.