Product Name: Recombinant Human IL-6RA (C-Fc)

Catalog #: PHH2285



Summary

Name IL-6 R alpha/IL-6RA/CD126/IL-6 Receptor Subunit alpha

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Interleukin-6 receptor subunit alpha is produced by our

Mammalian expression system and the target gene encoding Leu20-Pro365 is

expressed with a human IgG1 Fc tag at the C-terminus.

Accession # P08887

Host Human Cells

Species Human

Predicted Molecular Mass 65.4 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human IL-6RA (C-Fc)

Catalog #: PHH2285





Alternative Names

Interleukin-6 receptor subunit alpha; IL-6R subunit alpha; IL-6R-alpha; IL-6R 1; Membrane glycoprotein 80; gp80; CD126

Background

Interleukin 6 is a potent pleiotropic cytokine that regulates cell growth and differentiation and plays an important role in the immune response. IL6Ra is a part of the receptor for interleukin 6 cytokine. IL6Ra binds to IL6 with low affinity, but does not transduce a signal. Signal activation necessitates an association with IL6ST. Activation may lead to the regulation of the immune response, acute-phase reactions and hematopoiesis. Low concentration of a soluble form of IL6 receptor acts as an agonist of IL6 activity. Dysregulated production of IL6 and this receptor are implicated in the pathogenesis of many diseases, such as multiple myeloma, autoimmune diseases and prostate cancer.

Note

For Research Use Only, Not for Diagnostic Use.