

Product Name: Recombinant Human Neuropilin-1 (C-Fc)
Catalog #: PHH2329




Summary

Name	Neuropilin-1/NRP1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Neuropilin-1 is produced by our Mammalian expression system and the target gene encoding Phe22-Lys644 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH07533.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	96.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human Neuropilin-1 (C-Fc)
Catalog #: PHH2329



Alternative Names

CD304; NRP1; NRPNP1; VEGF165R; BDCA4

Background

Neuropilin-1 (Npn-1, previously neuropilin; also CD304) is a 130 - 140 kDa type I transmembrane (TM) glycoprotein that regulates axon guidance and angiogenesis. Two homologues, Neuropilin-1 and Neuropilin-2, are identified. Neuropilin-1 binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. NRP1 expression is regulated in EC by tumor necrosis factor- α , the transcription factors dHAND and Ets-1, and vascular injury. NRP1 upregulation is positively correlated with the progression of various tumors.

Note

For Research Use Only , Not for Diagnostic Use.