

**Product Name: Recombinant Human Neuropilin-1 (C-mFc)**  
**Catalog #: PHH2330**




## Summary

<b>Name</b>	Neuropilin-1/NRP1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Neuropilin-1 is produced by our Mammalian expression system and the target gene encoding Phe22-Lys644 is expressed with a mouse IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	AAH07533.1
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	96.2 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human Neuropilin-1 (C-mFc)**  
**Catalog #: PHH2330**



### Alternative Names

CD304; NRP1; NRPNP1; VEGF165R; BDCA4

### Background

Neuropilin-1 (Npn-1, previously neuropilin; also CD304) is a 130 - 140 kDa type I transmembrane (TM) glycoprotein that regulates axon guidance and angiogenesis. Two homologues, Neuropilin-1 and Neuropilin-2, are identified. Neuropilin-1 binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. NRP1 expression is regulated in EC by tumor necrosis factor- $\alpha$ , the transcription factors dHAND and Ets-1, and vascular injury. NRP1 upregulation is positively correlated with the progression of various tumors.

### Note

For Research Use Only , Not for Diagnostic Use.