


## Summary

Name	Adiponectin/Acrp30/ADIPOQ
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse Adipocyte Complement-related 30 kDa Protein is produced by our Mammalian expression system and the target gene encoding Glu18-Asn247 is expressed with a 6His tag at the C-terminus.
Accession #	Q60994
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	25.7 KDa
Formulation	Lyophilized from a 0.2 $\mu m$ filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}$ C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}$ C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## Background



Alternative NamesAdiponectin; 30 kDa Adipocyte Complement-Related Protein; Adipocyte<br/>complement-related 30 kDa protein; ACRP30; Adipocyte; C1q and Collagen<br/>Domain-Containing Protein; Adipose Most Abundant Gene Transcript 1 Protein;<br/>apM-1; Gelatin-Binding Protein; ADIPOQBackgroundAdiponectin is a secreted protein. It is synthesized exclusively by adipocytes and<br/>secreted into plasma. Adiponectin is an important adipokine that is involved in the<br/>control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-<br/>atherogenic and anti-inflammatory activities. Adiponectin Stimulates AMPK<br/>phosphorylation and activates in the liver and the skeletal muscle, enhancing<br/>glucose utilization and fatty-acid combustion. Adiponectin also antagonizes TNF-<br/>alpha by negatively regulating its expression in various tissues such as liver and<br/>macrophages, and also by counteracting its effects. It inhibits endothelial NF-

alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. It inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. Adiponectin may play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex: LMW, MMW or HMW.

## **Note** For Research Use Only , Not for Diagnostic Use.