

Product Name: Recombinant Mouse BMPR1A (C-Fc-6His)
Catalog #: PHM0041




Summary

Name	ALK-3/BMPR1A/BMPRIA
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Bone Morphogenetic Protein Receptor Type IA/Activin Receptor-like Kinase 3 is produced by our Mammalian expression system and the target gene encoding Gln24-Arg152 is expressed with a human IgG1 Fc, 6His tag at the C-terminus.
Accession #	P36895
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	42.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse BMPR1A (C-Fc-6His)
Catalog #: PHM0041



Alternative Names

ALK-3; Bone morphogenetic protein receptor type-1A; BMP type-1A receptor; BMPR-1A; Activin receptor-like kinase 3; BMP-2/BMP-4 receptor; Serine/threonine-protein kinase receptor R5; SKR5; CD292; Acvrlk3

Background

ALK-3 is a type I receptor for bone morphogenetic proteins (BMPs) which belong to the protein kinase superfamily, TKL Ser/Thr protein kinase family and TGF β receptor subfamily. The BMP receptors consist of the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. Seven known type I serine/threonine kinases and five mammalian type II serine/threonine kinase receptors function in TGF β superfamily signal transduction. The downstream molecules of the type I BMP receptors include the Smad (Smad1, 5 and 8) proteins that are phosphorylated in a ligand-dependent manner, and relay the BMP signal from the receptors to target genes in the nucleus. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. ALK-3 contains a GS domain and a protein kinase domain. ALK-3 is widely expressed. Defects in BMPR1A gene are a cause of a significant proportion of cases of Juvenile polyposis syndrome (JPS).

Note

For Research Use Only , Not for Diagnostic Use.