


Product Name: Recombinant Mouse BTNL2 (C-6His)
Catalog #: PHM0181



Summary

Name	BTNL2/Butyrophilin-like Protein 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Butyrophilin-like Protein 2 is produced by our Mammalian expression system and the target gene encoding Asp27-Ser452 is expressed with a 6His tag at the C-terminus.
Accession #	O70355
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	48.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse BTNL2 (C-6His)
Catalog #: PHM0181



Alternative Names

Butyrophilin-like protein 2; Btl2; Gm315; Ng9

Background

Butyrophilin-like 2 (BTNL2) is a member of the BTN/MOG Ig-superfamily and functions as a negative regulator of immune cell activation. Mouse BTNL2 is type I transmembrane glycoprotein that contains an extracellular domain (ECD), a transmembrane region and a short cytoplasmic domain. The ECD features two V-type Ig-like domains, two C-type Ig-like domains, and four glycosylation sites. BTNL2 is expressed in epithelial cells of the small intestine, colonic dendritic cells, and in cells of the lymph node. BTNL2 expression is upregulated in T cells following activation, a characteristic BTNL2 shares with the homologous B7 family of costimulatory molecules. BTNL2 negatively regulates T cells by inhibiting proliferation and inflammatory cytokine secretion. It also increases the expression of FoxP3 in T cells to promote regulatory T cell development. Single nucleotide polymorphisms in BTNL2 are associated with a risk for sporadic prostate cancer, rheumatoid arthritis, sarcoidosis, ulcerative colitis, and other inflammatory diseases.

Note

For Research Use Only , Not for Diagnostic Use.