
Product Name: Recombinant Mouse Carbonic Anhydrase 12 (C-Eiskilife Catalog #: PHM0210

Summary

Name	Carbonic Anhydrase 12/CA12/CA-XII
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse Carbonic Anhydrase 12 is produced by our Mammalian expression system and the target gene encoding Ala25-Ser301 is expressed with a 6His tag at the C-terminus.
Accession #	Q8Cl85
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	32.4 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative NamesCarbonic anhydrase 12; Carbonate dehydratase XII; Carbonic anhydrase XII; CA-XII;
CA12; Carbonate dehydratase XII; CAXIIBackgroundCarbonic Anhydrase (CA) XII, also known as Car12 and CA12, is an extracellular
enzyme involved in the regulation of the microenvironment acidity and tumor
malignant phenotype, was originally identified as a protein overexpressed in some
types of cancers. It has showed that CA XII is induced by hypoxia and oestrogen
and expressed at high levels on various types of cancer. The enzyme is directly
involved in tumour progression, and its inhibition has an anti-tumour effect. Apart
from its role in carcinogenesis, the enzyme contributes to various other diseases
like glaucoma and arteriosclerotic plaques, among others. CA XII is therefore
regarded as promising target for specific therapies, and may be used as a novel
prognostic marker in combination with histologic grade of the tumors.

Note

For Research Use Only, Not for Diagnostic Use.