

**Product Name: Recombinant Mouse Carbonic Anhydrase 12 (C-6His)**  
**Catalog #: PHM0210**




## Summary

<b>Name</b>	Carbonic Anhydrase 12/CA12/CA-XII
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse Carbonic Anhydrase 12 is produced by our Mammalian expression system and the target gene encoding Ala25-Ser301 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	Q8CI85
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	32.4 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Mouse Carbonic Anhydrase 12 (C-6 His)**  
**Catalog #: PHM0210**



### Alternative Names

Carbonic anhydrase 12; Carbonate dehydratase XII; Carbonic anhydrase XII; CA-XII; CA12; Carbonate dehydratase XII; CAXII

### Background

Carbonic Anhydrase (CA) XII, also known as Car12 and CA12, is an extracellular enzyme involved in the regulation of the microenvironment acidity and tumor malignant phenotype, was originally identified as a protein overexpressed in some types of cancers. It has showed that CA XII is induced by hypoxia and oestrogen and expressed at high levels on various types of cancer. The enzyme is directly involved in tumour progression, and its inhibition has an anti-tumour effect. Apart from its role in carcinogenesis, the enzyme contributes to various other diseases like glaucoma and arteriosclerotic plaques, among others. CA XII is therefore regarded as promising target for specific therapies, and may be used as a novel prognostic marker in combination with histologic grade of the tumors.

### Note

For Research Use Only , Not for Diagnostic Use.