# Product Name: Recombinant Mouse Nectin-2 (C-6His) Catalog #: PHM0283



### **Summary**

Name Nectin-2/CD112/PVRL2/Pvr/Mph/Pvs

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Poliovirus Receptor-Related Protein 2 is produced by

our Mammalian expression system and the target gene encoding Gln32-

Gly351 is expressed with a 6His tag at the C-terminus.

Accession # P32507

**Host** Human Cells

**Species** Mouse

Predicted Molecular Mass 35.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## **Background**

# Product Name: Recombinant Mouse Nectin-2 (C-6His) Catalog #: PHM0283



Alternative Names CD112; n

CD112; nectin2; Herpes virus entry mediator B; Herpesvirus entry mediator B; HveB;Murine herpes virus entry protein B; mHveB;Poliovirus receptor homolog;

Poliovirus receptor-related protein 2; Pvrl2

Background

Nectin-2( CD112) is a member of the nectin family, which contains two Ig-like C2-type domains and one Ig-like V-type domain in the extracellular region. Nectins are type I transmembrane glycoproteins that are calcium-independent immunoglobulin (Ig)-like cell adhesion molecules (CAMs). Nectin2 is widely expressed in human tissues including brain, spinal cord, spleen, kidney, heart and liver. It can form trans-heterodimers with PVRL3/nectin-3 and interacts with CD226. Mutations of alleles of the murine CD112 gene can result in conditions such as morphologically aberrant spermatozoa. It may function in allergic reactions, and accordingly may used as a novel target for anti-allergic therapy.

#### Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838