

Product Name: Recombinant Mouse CD27 (C-Fc-6His)
Catalog #: PHM0321




Summary

Name	CD27/TNFRSF7
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse CD27 Antigen is produced by our Mammalian expression system and the target gene encoding Pro24-Arg182 is expressed with a human IgG1 Fc, 6His tag at the C-terminus.
Accession #	P41272
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	45.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse CD27 (C-Fc-6His)
Catalog #: PHM0321



Alternative Names

CD27 antigen; CD27L receptor; T-cell activation antigen CD27; Tumor necrosis factor receptor superfamily member 7; CD27

Background

Mouse CD27 is a single-pass type I membrane protein which is a member of the tumor necrosis factor receptor superfamily. CD27 is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. CD27 transduces signals that lead to the activation of NF- κ B and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Note

For Research Use Only , Not for Diagnostic Use.