# **Product Name: Recombinant Mouse CPA2 (C-6His)**

Catalog #: PHM0449



### **Summary**

Name CPA2/Carboxypeptidase A2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Carboxypeptidase A2 is produced by our Mammalian

expression system and the target gene encoding Gln17-Tyr417 is expressed

with a 6His tag at the C-terminus.

Accession # Q504N0

**Host** Human Cells

**Species** Mouse

Predicted Molecular Mass 46.2 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


## **SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

## **Product Name: Recombinant Mouse CPA2 (C-6His)**

Catalog #: PHM0449





#### **Alternative Names**

CPA2; Carboxypeptidase A2

### **Background**

Mouse carboxypeptidase A2(CPA2) is a secreted pancreatic procarboxy -peptidase which belongs to the peptidase M14 family. CPA2 consists of a signal peptide, a pro region and a mature chain. It can be activated after cleavage of the pro peptide. CPA2 cleaves the C-terminal amide or ester bond of peptides that have a free C-terminal carboxyl group. The hydrolytic action of CPA2 was identified with a preference towards long substrates with aromatic amino acids in their C-terminal end, particularly tryptophan.

#### Note

For Research Use Only, Not for Diagnostic Use.