

Product Name: Recombinant Mouse Decorin (C-6His)
Catalog #: PHM0524




Summary

Name	Decorin/PGS2/PG40
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Decorin is produced by our Mammalian expression system and the target gene encoding Gly17-Lys354 is expressed with a 6His tag at the C-terminus.
Accession #	P28654
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	39 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse Decorin (C-6His)
Catalog #: PHM0524



Alternative Names

Decorin; Bone proteoglycan II; PG-S2; PG40; Dcn

Background

Decorin, also known as PG40 and DCN, is a member of the class I family of small leucine-rich proteoglycans (SLRPs) that is expressed in the stroma of various forms of cancer and has been recently proposed to act as a guardian from the matrix. Mature human Decorin contains 12 tandem LRR and shares 80% and 78% aa sequence identity with mouse and rat Decorin, respectively. Decorin embraces numerous functions including: regulation of collagen fibrillogenesis, hepatic carcinogenesis, fetal membrane and calcium homeostasis, keratinocyte function, and suppression of angiogenesis. Most recently, soluble decorin has been shown to induce autophagy in endothelial cells and mitophagy in breast carcinoma cells.

Note

For Research Use Only , Not for Diagnostic Use.