Product Name: Recombinant Mouse CD39 (C-6His)

Catalog #: PHM0574



Summary

Name CD39/ENTPD1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Ectonucleoside Triphosphate Diphosphohydrolase 1 is

produced by our Mammalian expression system and the target gene

encoding Thr38-Ile478 is expressed with a 6His tag at the C-terminus.

Accession # P55772

Host Human Cells

Species Mouse

Predicted Molecular Mass 50.5 KDa

Formulation Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 500mM NaCl, 10%

Glycerol, pH 7.4.

Shipping The product is shipped on dry ice/polar packs. Upon receipt, store it immediately

at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Mouse CD39 (C-6His)

Catalog #: PHM0574





Alternative Names

Ectonucleoside triphosphate diphosphohydrolase 1; NTPDase 1; Ecto-ATP diphosphohydrolase 1; Ecto-ATPDase 1; Ecto-ATPase 1; Ecto-apyrase; Lymphoid cell activation antigen; CD39

Background

Ectonucleoside triphosphate diphosphohydrolase-1(NTPDase-1) is an integral membrane protein with an extracellular active site. NTPDase-1 was originally described as CD39, a B lymphocyte cell surface marker. But it is also present on the surface of natural killer cells, T cells, and some endothelial cells. NTPDase1 hydrolyzes the β-andy phosphate residues of nucleotides, preferring ATP as the substrate. Through its hydrolysis of extracellular nucleotides, NTPDase-1 plays a role in the regulation of purinergic signaling. NTPDase-1 is involved in the processes of thromboregulation and vascular inflammation. The administration of soluble NTPDase-1 may have therapeutic applications for the treatment of some vascular and transplantation-associated diseases.

Note

For Research Use Only, Not for Diagnostic Use.