Product Name: Recombinant Mouse EFNA5 (C-6His)

Catalog #: PHM0593



Summary

Name Ephrin-A5/EFNA5

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Ephrin-A5 is produced by our Mammalian expression

system and the target gene encoding Gln21-Gln206 is expressed with a 6His

tag at the C-terminus.

Accession # 008543

Host Human Cells

Species Mouse

Predicted Molecular Mass 22.5 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Mouse EFNA5 (C-6His)

Catalog #: PHM0593





Alternative Names

Ephrin-A5; AL-1; EPH-related receptor tyrosine kinase ligand 7; Epl7; Eplg7; Lerk7; Efna5

Background

Ephrin-A5 is a glycosylphosphatidylinositol (GPI)-anchored protein of the ephrin-A subclass of ephrin ligands that binds to the EphA subclass of Eph receptors. Ephrin-A5 has also been shown to bind to the EphB2 receptor. It is crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Ephrin-A5 binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling.

Note

For Research Use Only, Not for Diagnostic Use.