Product Name: Recombinant Mouse FcRL1 (C-6His)

Catalog #: PHM0627



Summary

Name FCRL1/FcRH1/CD307a/Fc Receptor-like Protein 1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Fc Receptor-like Protein 1 is produced by our

Mammalian expression system and the target gene encoding Ala17-Ser219 is

expressed with a 6His tag at the C-terminus.

Accession # BAC30017.1

Host Human Cells

Species Mouse

Predicted Molecular Mass 22.5 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse FcRL1 (C-6His) Catalog #: PHM0627



Alternative Names Fc receptor-like protein 1; FcR-like protein 1; FcRL1; BXMAS1-like protein 1;

mBXMH1; Fc receptor homolog 1; FcRH1; moFcRH1; IFGP family protein 1;

mIFGP1; CD307a

Background Mouse Fc receptor-like protein 1 (FCRL1) is a single-pass type I membrane protein.

It is expressed in B-cells at the various stages of differentiation. Mouse FCRL1 consists of a 203 amino acid (aa) extracellular domain (ECD) with two Ig-like domains, a 21 aa transmembrane segment, and a 103 aa cytoplasmic domain with two immunotyrosine activation motifs (ITAMs). FCRL1 is likely not a receptor for immunoglobulin. It may function as an activating coreceptor in B-cells and B-cells

differentiation.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838