

Product Name: Recombinant Mouse Asprosin (N-8His)
Catalog #: PHM0655




Summary

Name	Asprosin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Fibrillin-1 is produced by our Mammalian expression system and the target gene encoding Ser2732-His2871 is expressed with a 8His tag at the N-terminus.
Accession #	AAA56840.1
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	16.9 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse Asprosin (N-8His)
Catalog #: PHM0655



Alternative Names

Fibrillin-1; Fbn1; Asprosin; Fbn-1

Background

Asprosin is a protein hormone that is produced by white adipose tissue in mammals (and potentially by other tissues), which is then transported to the liver and stimulates it to release glucose into the blood stream. In the liver asprosin activates rapid glucose release by a cAMP-dependent pathway. The glucose release by the liver into the blood stream is vital for brain function and survival during fasting. People with neonatal progeroid syndrome lack asprosin, while people with insulin resistance have it in abundance. In animal tests asprosin showed potential for treating type 2 diabetes. When antibodies targeting asprosin were injected into diabetic mice, blood glucose and insulin levels improved.

Note

For Research Use Only , Not for Diagnostic Use.